

Discrete Mathematics 103 (1992) 57–65
North-Holland

57

A minimax theorem for infinite graphs with ideal points

Norbert Polat

*Département de Mathématiques, Bâtiment Doyen Jean Braconnier, Université Claude Bernard (Lyon I), 43, Boulevard du 11 Novembre 1918, 69622 Villeurbanne Cedex, France
I.A.E., Université Jean Moulin (Lyon III), 15, quai Claude Bernard, 69239 Lyon Cedex 2, France*

Received 10 April 1990

Abstract

Polat, N., A minimax theorem for infinite graphs with ideal points, *Discrete Mathematics* 103 (1992) 57–65.

Let \mathcal{A} be a family of sets of ends of an infinite graph, having the property that every element of any member of \mathcal{A} can be separated from the union of all other members by a finite set of vertices. By defining appropriate concepts of \mathcal{A} -paths and of \mathcal{A} -separators, we show that there are a set of pairwise disjoint \mathcal{A} -paths and an \mathcal{A} -separator which have the same 'cardinality'.

Introduction

In a recent paper [8] we generalized different extensions [10, 3, 5] of Menger's Theorem to infinite graphs admitting their ends as ideal points. We proved, by defining appropriate concepts of AB -paths and of AB -separators, that, if A and B are two sets of ends of an infinite graph G , having the property that any element of A (resp. B) can be separated from B (resp. A) by a finite set of vertices, then there are a set of pairwise disjoint AB -paths and an AB -separator which have the same cardinality.

In this result, as in any Menger-like theorem, the separation of exactly two sets of objects of a graph is involved. A more interesting and much more general result would be a similar theorem dealing with any set $\mathcal{A} := \{A_i : i \in I\}$ of pairwise disjoint sets of ends of a graph G . This is this generalization which is studied in this paper.

For such a set $\mathcal{A} = \{A_i : i \in I\}$ we define an \mathcal{A} -path as an $A_i A_j$ -path (as defined in [8]) for any different $i, j \in I$, and an \mathcal{A} -separator as an ordered pair (S, T) with $S \subseteq V(G)$ and $T \subseteq E(G)$ such that any \mathcal{A} -path has a vertex in S or an edge in T , and whose deletion does not split or destroy any end in $\bigcup \mathcal{A}$. The existence of an \mathcal{A} -separator is ensured when, for any $i \in I$, every element of A_i can be

separated from $\bigcup_{j \neq i} A_j$ by a finite set of vertices. Then, using the extension to infinite graphs of a theorem of Mader [4] that we give in Section 1, we get a minimax theorem between the cardinalities of the pairwise disjoint sets of \mathcal{A} -paths and those—actually a modified version of cardinality—of the \mathcal{A} -separators, for any set \mathcal{A} having the preceding property, so generalizing Theorem 3.3 of [8].

0. Preliminaries

0.1. If X is a set we denote by $|X|$ its cardinality, by $\mathcal{P}(X)$ its power set, and if n is a cardinal, by $[X]^n$ (resp. $[X]^{<n}$, $[X]^{\geq n}$) the set of its subsets of cardinality n (resp. $<n$, $\geq n$). Furthermore, $\lfloor |X|/2 \rfloor$ will be $\max\{n \in \mathbb{N} : n \leq |X|/2\}$ if X is finite, and $|X|$ otherwise.

0.2. A graph G is a set $V(G)$ (*vertex set*) together with a set $E(G) \subseteq [V(G)]^2$ (*edge set*). H is a *subgraph* of G ($H \subseteq G$) if $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$. H is an *induced subgraph* if $E(H) = [V(H)]^2 \cap E(G)$. For $A \subseteq V(G)$ we denote by $G \upharpoonright A$ the subgraph induced by A . If B is any set, and H any graph, we define $G - B := G \upharpoonright (V(G) - B)$ and $G - H := G - V(H)$. For $T \subseteq E(G)$ we denote by $G(T)$ the subgraph of G whose edge set is T , and vertex set is $\bigcup T$. The *union* of a family $(G_i)_{i \in I}$ of graphs is the graph $G := \bigcup_{i \in I} G_i$ given by $V(G) = \bigcup_{i \in I} V(G_i)$ and $E(G) = \bigcup_{i \in I} E(G_i)$. The *intersection* is defined similarly. If X and Y are subgraphs of G , then the *boundary of X with Y* is the set $\mathcal{B}(X, Y) := \{x \in V(X) : \{x, y\} \in E(G) \text{ for some } y \in V(Y - X)\}$. In particular $\mathcal{B}(X) := \mathcal{B}(X, G)$ is the *boundary* of X . The set of components of G is denoted by \mathcal{C}_G , and if x is a vertex, then $\mathcal{C}_G(x)$ is the component of G containing x . A *path* $W := \langle x_0, \dots, x_n \rangle$ is a graph with $V(W) = \{x_0, \dots, x_n\}$, $x_i \neq x_j$ if $i \neq j$, and $E(W) = \{\{x_i, x_{i+1}\} : 0 \leq i < n\}$. A *ray* or *one-way infinite path* $R := \langle x_0, x_1, \dots \rangle$, and a *double ray* or *two-way infinite path* $D := \langle \dots, x_{-1}, x_0, x_1, \dots \rangle$ are defined similarly. A path $\langle x_0, \dots, x_n \rangle$ is called an $x_0 x_n$ -*path*. For $A, B \subseteq V(G)$, an AB -*path* of G , or simply an A -*path* when $A = B$, is an xy -path of G whose only vertices in $A \cup B$ are x and y , with $x \in A$ and $y \in B$. Two paths are *internally disjoint* if they have at most their endpoints in common.

0.3. The *ends* of a graph G (this concept was introduced by Freudenthal [1] and independently by Halin [2]) are the classes of the equivalence relation \sim_G defined on the set of all rays of G by $R \sim_G R'$ if and only if there is a ray R'' whose intersections with R and R' are infinite; or equivalently if and only if $\mathcal{C}_{G-S}(R) = \mathcal{C}_{G-S}(R')$ for any $S \in [V(G)]^{<\omega}$ (where $\mathcal{C}_{G-S}(R)$ denotes the component of $G - S$ containing a subray of R). We will use the following notations:

- if R is a ray of G , then $[R]_G$ is the class of R modulo \sim_G ;
- $\mathfrak{T}(G)$ is the set of all ends of G ;
- if H is a subgraph of G , then $\mathfrak{T}_H(G)$ is the set of ends of G having rays of H as elements;
- if $\tau \in \mathfrak{T}(G)$ and $S \subseteq V(G)$, then $\mathcal{C}_{G-S}(\tau)$ is the component of $G - S$ containing a ray in τ ; and if $A \subseteq \mathfrak{T}(G)$ then $\mathcal{C}_{G-S}(A) := \bigcup_{\tau \in A} \mathcal{C}_{G-S}(\tau)$;
- if $\tau \in \mathfrak{T}(G)$ then $V_\tau := \{x \in V(G) : x \in V(\mathcal{C}_{G-S}(\tau)) \text{ for any } S \in [V(G) - \{x\}]^{<\omega}\}$.

0.4. For every $S \in [V(G)]^{<\omega}$ we denote by S^* the equivalence relation on $\mathfrak{T}(G)$ defined by $(\tau, \tau') \in S^*$ if and only if $\mathcal{C}_{G-S}(\tau) = \mathcal{C}_{G-S}(\tau')$. The family of all such equivalence relations is a base of a Hausdorff uniformity on $\mathfrak{T}(G)$ (see [5]). We will assume that, throughout this paper, the end set $\mathfrak{T}(G)$ of a graph G is endowed with the topology induced by this uniformity, i.e., the topology such that, for every end τ , the family $(S^*(\tau))_{S \in [V(G)]^{<\omega}}$ is a local basis at the point τ . For any $A \subseteq \mathfrak{T}(G)$ we will denote by \bar{A} the closure of A for this topology.

0.5. An infinite subset S of $V(G)$ is *concentrated* in G if it has the following equivalent properties [5, Theorem 3.5]:

- (i) there is an end τ such that $S - V(\mathcal{C}_{G-F}(\tau))$ is finite for any $F \in [V(G)]^{<\omega}$ (S is said to be *concentrated in τ*);
- (ii) for all $T, U \in [S]^{>\omega}$, there is a family of pairwise TU -paths in G .

0.6. Let $A \subseteq \mathfrak{T}(G)$. A subset S of $V(G)$ is *A-dispersed* if it has the following equivalent properties [8, Proposition 0.6]:

- (i) for every $\tau \in A$ there is an $F \in [V(G)]^{<\omega}$ such that $S \cap V(\mathcal{C}_{G-F}(\tau)) = \emptyset$;
- (ii) S has no subset concentrated in some end $\tau \in A$.

We denote by $D(A)$ the set of A -dispersed subsets of $V(G)$. $S \in D(A)$ is *finitary* if, for any $\tau \in A$, there is a finite subset F of S such that $\mathcal{C}_{G-F}(\tau) = \mathcal{C}_{G-S}(\tau)$. By Proposition 0.13 of [8], for any $S \in D(A)$ there is a finitary $T \in D(A)$ which contains S .

1. An extension of a theorem of Mader to infinite graphs

1.1 Let A be an independent set of vertices of a graph G (i.e., $E(G) \cap [A]^2 = \emptyset$). An ordered pair (S, T) is called an *A-separator* of G if it has the following properties:

- (i) $S \subseteq V(G) - A$;
- (ii) $T \subseteq E(G) \cap [V(G) - A \cup S]^2$;
- (iii) for every A -path W , $V(W) \cap S \neq \emptyset$ or $E(W) \cap T \neq \emptyset$.

We will denote by $\text{DISJ}_G(A)$ the set of all sets of pairwise internally disjoint A -paths of G , and by $\text{SEP}_G(A)$ the set of A -separators of G ; and we set

$$\text{disj}_G(A) := \sup\{|\mathcal{D}| : \mathcal{D} \in \text{DISJ}_G(A)\}$$

and

$$\text{sep}_G(A) := \inf\{|(S, T)| : (S, T) \in \text{SEP}_G(A)\}$$

where

$$|(S, T)| := |S| + \sum_{X \in \mathcal{C}_{G(T)}} \lfloor |\mathcal{B}(X, G - S)|/2 \rfloor.$$

When no confusion is likely we will write $\text{DISJ}(A)$, $\text{SEP}(A)$, $\text{disj}(A)$ and $\text{sep}(A)$. Mader proved in [4] that $\text{disj}_G(A) = \text{sep}_G(A)$ for any finite graph G . We will now extend this result to arbitrary infinite graphs.

Theorem 1.2. *Let A be an independent set of vertices of a graph G . Then $\text{disj}(A) = \text{sep}(A)$.*

Proof. We have clearly $\text{disj}(A) \leq \text{sep}(A)$. If G is finite, then the converse is Mader's Theorem. Assume that G is infinite. We have two cases:

Case 1: $\text{disj}(A) \geq \omega$.

Let \mathcal{D} be an element of $\text{Disj}(A)$ which is maximal with respect to inclusion. Then $(V(\bigcup \mathcal{D}), \emptyset)$ is an A -separator of cardinality $|\mathcal{D}|$ since all A -paths are finite. Hence $\text{disj}(A) = \text{sep}(A)$.

Case 2: $\text{disj}(A) < \omega$.

This is the nontrivial case. Let $\mathcal{D} \in \text{DISJ}(A)$ be of maximum cardinality $n := \text{disj}(A)$. Suppose that $\text{sep}(A) > n$. Let \mathcal{H} be the set of finite induced subgraphs of G containing \mathcal{D} , and let $H \in \mathcal{H}$. By Mader's Theorem $\text{disj}_H(A_H) = \text{sep}_H(A_H) = n$, where $A_H = A \cap V(H)$. Let $(S, T) \in \text{SEP}_H(A_H)$ be such that $|(S, T)| = n$. For any $W \in \mathcal{D}$ and any $X \in \mathcal{C}_{H(T)}$ with $E(X) \cap E(W) \neq \emptyset$, we have $|\mathcal{B}(X) \cap V(W)| \geq 2$. Thus, since $|(S, T)| = n$, the set of elements of \mathcal{D} meeting S has the same cardinality as S . Hence, since $V(\bigcup \mathcal{D})$ is finite, there is $S \subseteq V(\bigcup \mathcal{D})$ with $|S \cap V(W)| \leq 1$ for every $W \in \mathcal{D}$, such that, for any $H \in \mathcal{H}$, there is $T_H \subseteq E(H)$ with $(S, T_H) \in \text{SEP}_H(A_H)$ and $|(S, T_H)| = n$.

For any $H \in \mathcal{H}$ we define the function $\theta_H \in \prod_{e \in E(H)} \{0, 1\}$ so that, if $T_H := \theta_H^{-1}(1)$, then $(S, T_H) \in \text{SEP}_H(A_H)$ with $|(S, T_H)| = n$. W.l.o.g. we can suppose that T_H is such that each component of $G(T_H)$ is an induced subgraph of H , thus of G . By Rado's Selection Principle [9], there is a function $\theta \in \prod_{e \in E(G)} \{0, 1\}$ such that, for any $H \in \mathcal{H}$, there is $H' \in \mathcal{H}$ with $H \subseteq H'$ and $\theta|_{E(H)} = \theta_{H'}|_{E(H)}$. Let $T := \theta^{-1}(1)$.

Let Y be any finite subgraph of $G \setminus T$, and let W be an A -path of G . There is $H \in \mathcal{H}$ such that $Y \cup W \subseteq H$ and $\theta|_{E(Y \cup W \cup \bigcup \mathcal{D})} = \theta_H|_{E(Y \cup W \cup \bigcup \mathcal{D})}$. Then, on the one hand, Y is a subgraph of $H \setminus T_H$, and this proves that $A \cup S$ does not meet $V(Y)$, thus $\bigcup T$; so T satisfies condition (ii) of Definition 1.1. On

the other hand $S \cap V(W) = \emptyset$ implies $T \cap E(W) = T_H \cap E(W) \neq \emptyset$. Thus (S, T) is an A -separator of G .

We will show that $|(S, T)| = n$. Suppose that $|(S, T)| > n$. Then there exist $X_1, \dots, X_p \in \mathcal{C}_{G(T)}$ with $X_i \neq X_j$ if $i \neq j$, and for $1 \leq i \leq p$ there is a finite subset B_i of $\mathcal{B}(X_i, G - S)$, such that

$$\sum_{1 \leq i \leq p} [|B_i|/2] > n - |S|.$$

For $1 \leq i \leq p$ there is a finite connected induced subgraph Y_i of X_i containing B_i . Let $Y'_i := Y_i \cup \{x, y_x\} : x \in B_i, y_x \text{ is a vertex of } G - X_i \text{ such that } \{x, y_x\} \in E(G) - T$. Since p , as well as each Y'_i , is finite, there is $H \in \mathcal{H}$ containing $Y := \bigcup_{1 \leq i \leq p} Y'_i$ such that $\theta_H |E(Y \cup \bigcup \mathcal{D}) = \theta |E(Y \cup \bigcup \mathcal{D})$. Then for $1 \leq i \leq p$, on the one hand Y_i is a connected subgraph of a component Z_i of $H(T_H)$, and on the other hand $\{x, y_x\} \notin T_H$ for every $x \in B_i$. Thus $B_i \subseteq \mathcal{B}_H(Z_i, H - S)$ —where \mathcal{B}_H denotes the boundary in the graph H —since, by the assumption we made, each Y'_i is an induced subgraph of H . Let $\mathcal{Z} := \{Z_1, \dots, Z_p\}$; notice that we may have $Z_i = Z_j$ for some $i \neq j$. Then

$$\sum_{Z \in \mathcal{Z}} [|\mathcal{B}_H(Z, H - S)|/2] \geq \sum_{1 \leq i \leq p} [|B_i|/2] > n - |S|,$$

which is a contradiction with

$$\sum_{Z \in \mathcal{Z}} [|\mathcal{B}_H(Z, H - S)|/2] \leq \sum_{Z \in \mathcal{C}_{H(T_H)}} [|\mathcal{B}_H(Z, H - S)|/2] = n - |S|.$$

hence $|(S, T)| = n$. Therefore $\text{disj}_G(A) = \text{sep}_G(A)$. \square

2. \mathcal{A} -separators

From now on, $\mathcal{A} := \{A_i : i \in I\}$ will be a set of pairwise disjoint subsets of the end set $\mathfrak{I}(G)$ of some infinite graph G .

Definition 2.1. An ordered pair $(S, T) \in \mathcal{P}(V(G)) \times \mathcal{P}(E(G))$ is an \mathcal{A} -separator if it has the following properties:

- (i) $S \cap \bigcup T = \emptyset$;
- (ii) $S \cup \bigcup T \in D(\bigcup \mathcal{A})$;
- (iii) for every double ray W consisting of two rays $R_i \in \bigcup \mathcal{A}_i$ and $R_j \in \bigcup \mathcal{A}_j$ with $\{i, j\} \in [I]^2$, $V(W) \cap S \neq \emptyset$ or $E(W) \cap T \neq \emptyset$.

We will denote by $\text{SEP}(\mathcal{A})$ the set of \mathcal{A} -separators and we will set

$$\text{sep}(\mathcal{A}) := \inf\{|(S, T)| : (S, T) \in \text{SEP}(\mathcal{A})\}$$

where $|(S, T)|$ is defined as in Definition 1.1.

Remarks 2.2. We have clearly:

(2.2.1) $(S, T) \in \text{SEP}(\mathcal{A})$ if and only if $(S, T) \in \text{SEP}(A_i, A_j)$ for every $\{i, j\} \in [I]^2$.

(2.2.2) If $(S, T) \in \text{SEP}(\mathcal{A})$ then $(S \cup T, \emptyset) \in \text{SEP}(\mathcal{A})$.

(2.2.3) $(S, \emptyset) \in \text{SEP}(\mathcal{A})$ if and only if $\mathcal{G}_{G-S}(A_i) \cap \mathcal{C}_{G-S}(A_j) = \emptyset$ for every $\{i, j\} \in [I]^2$.

If (S, \emptyset) is an \mathcal{A} -separator, we will say simply that S is an \mathcal{A} -separator, and we will write $S \in \text{SEP}(\mathcal{A})$.

Theorem 2.3. $\text{SEP}(\mathcal{A}) \neq \emptyset$ if and only if $A_i \cap \overline{\bigcup_{j \neq i} A_j} = \emptyset$ for every $i \in I$.

Proof. In the following, if $(X_\xi)_{\xi < \alpha}$ is any sequence we will denote $\bigcup_{\xi < \alpha} X_\xi$ by $X_{(\alpha)}$.

(a) Assume that $\mathcal{A} = \{A_\xi : \xi < \alpha\}$ where α is an ordinal, with $A_\xi \neq A_\eta$ if $\xi \neq \eta$, and suppose that $A_\xi \cap \bigcup_{\eta \neq \xi} A_\eta = \emptyset$ for every $\xi < \alpha$. We define by induction a sequence $(S_\xi)_{\xi < \alpha}$ such that, for any $\xi < \alpha$:

(i) $S_{(\xi)} \in D(\bigcup \mathcal{A})$ with $S_{(0)} := \emptyset$;

(ii) $S_\xi \in D(\bigcup \mathcal{A}) \cap \mathcal{P}(V(G) - S_{(\xi)})$ such that $S_{(\xi+1)} \in \text{SEP}(A_\xi, \bigcup \mathcal{A} - A_\xi)$, and S_ξ is minimal for this property with respect to inclusion.

Suppose that $S_{(\beta)}$ is defined for some $\beta < \alpha$, such that $S_{(\beta)} \in D(\bigcup \mathcal{A})$. Let $G_\beta := G - S_{(\beta)}$, and for each $\xi < \alpha$ let $A_\xi^\beta := \{[R]_{G_\beta} : R \text{ is a ray of } G_\beta \text{ with } [R]_G \in A_\xi\}$. The set $\mathcal{A}^\beta := \{A_\xi^\beta : \xi < \alpha\}$ has the same properties as \mathcal{A} , since $S_{(\beta)}$ is $\bigcup \mathcal{A}$ -dispersed. By Corollary 2.8 of [7], there is a connected induced subgraph H_β of G_β with $\mathfrak{T}_{H_\beta}(G_\beta) = \bigcup_{\beta < \xi < \alpha} A_\xi^\beta$ and such that the boundary of H_β with any component of $G_\beta - H_\beta$ is finite. Let

$$S_\beta := \bigcup_{\tau \in A_\beta^\beta} \mathcal{B}(H_\beta, \mathcal{C}_{G_\beta - H_\beta}(\tau)).$$

Obviously $S_\beta \in D(A_\beta^\beta) \subseteq D(A_\beta)$. Suppose $S_\beta \notin D(\bigcup \mathcal{A} - A_\beta)$. Then $S_\beta \notin D(\bigcup \mathcal{A}^\beta - A_\beta^\beta)$. Thus there is an infinite subset of S_β which is concentrated in some end $\tau \in A_\xi^\beta$ for some $\xi \neq \beta$. But since $\mathcal{B}(H_\beta, \mathcal{C}_{G_\beta - H_\beta}(\tau'))$ is finite for every $\tau' \in A_\beta^\beta$, we have $\tau \in A_\beta^\beta$, a contradiction with $A_\xi^\beta \cap A_\beta^\beta = \emptyset$. Thus $S_\beta \in D(\mathcal{A})$. Therefore clearly $S_{(\beta+1)} \in \text{SEP}(\mathcal{A}_\beta, \bigcup \mathcal{A} - A_\beta)$, since

$$\mathfrak{T}_{H_\beta}(G_\beta) = \overline{\bigcup_{\beta < \xi < \alpha} A_\xi^\beta},$$

and if $\beta > 0$, $S_{(\beta)} \in \text{SEP}(A_\xi, \bigcup \mathcal{A} - A_\xi)$ for every $\xi < \beta$. Furthermore, by its definition as a boundary, S_β is minimal with respect to inclusion.

It remains to prove that $S_{(\beta)} \in D(\bigcup \mathcal{A})$ for any $\beta < \alpha$. This is clear if β is a successor ordinal, by condition (i) and by the fact that $D(\bigcup \mathcal{A})$ is closed under finite unions. Assume that β is a limit ordinal, and suppose that $S_{(\beta)}$ has an infinite subset which is concentrated in some end θ . For any $\xi < \beta$, by the minimality of S_η for all $\eta < \beta$ (condition (ii)), we have $\mathcal{C}_{G - S_{(\eta)}}(\tau) = \mathcal{C}_{G - S_{(\xi)}}(\tau)$ for every $\tau \in A_\xi$. Thus $\theta \notin A_{(\beta)}$. Suppose $\theta \notin A_{(\beta)}$. Then there is a finite subset F of $V(G - \mathcal{C}_{G - S_{(\beta)}}(A_{(\beta)}))$ such that $F \in \text{SEP}(\theta, A_{(\beta)})$. But since, by condition (ii), the

S_ξ 's are minimal with respect to inclusion, every element of $S_{(\beta)}$ is adjacent to a vertex of $\mathcal{C}_{G-S_{(\beta)}}(A_{(\beta)})$. Hence, by the definition of F , we must have $\overline{S_{(\beta)}} \cap V(\mathcal{C}_{G-F}(\theta)) = \emptyset$, a contradiction with the definition of θ . Therefore $\theta \in A_{(\beta)} - A_{(\beta)}$; and this proves that $\theta \notin \bigcup \mathcal{A}$, thus that $S_{(\beta)}$ is $\bigcup \mathcal{A}$ -dispersed. Finally $S_{(\alpha)} \in \text{SEP}(\mathcal{A})$.

(b) Conversely suppose that $A_i \cap \overline{\bigcup_{j \neq i} A_j} \neq \emptyset$ for some $i \in I$, and let τ be an element of this intersection. If $S \in D(\bigcup \mathcal{A})$ then $S \cap V(\mathcal{C}_{G-F}(\tau)) = \emptyset$ for some finite subset F of $V(G)$. But $\tau \in \overline{\bigcup_{j \neq i} A_j}$ implies that $\mathcal{C}_{G-F}(\tau) = \mathcal{C}_{G-F}(\tau')$ for some $\tau' \in \bigcup_{j \neq i} A_j$. Hence $S \notin \text{SEP}(A_i, \bigcup_{j \neq i} A_j) \supseteq \text{SEP}(\mathcal{A})$. \square

In particular $\text{SEP}([\mathfrak{X}(G)]^1) \neq \emptyset$ if and only if the space $\mathfrak{X}(G)$ is discrete, that is if every end of G is a free end.

2.4. For $(S, T), (S', T') \in \text{SEP}(\mathcal{A})$ we set $(S, T) \leq (S', T')$ if and only if $S \subseteq S'$ and $T \subseteq T'$. This relation is obviously an order on $\text{SEP}(\mathcal{A})$. We will show that \mathcal{A} has minimal separators with respect to \leq .

Proposition 2.5. *The poset $(\text{SEP}(\mathcal{A}), \geq)$ is inductive.*

Proof. Let $\mathcal{C} := \{(S_k, T_k) : k \in K\}$ be a decreasing chain of $\text{SEP}(\mathcal{A})$, and let $S := \bigcap_{k \in K} S_k$ and $T := \bigcap_{k \in K} T_k$. Conditions (i) and (ii) of definition 2.1 are clearly satisfied by (S, T) . Let W be a double ray consisting of a ray $W_i \in \bigcup A_i$ and a ray $W_j \in \bigcup A_j$ for some pair $\{i, j\}$ of distinct elements of I . For any $k \in K$, the set $V(W) \cap (S_k \cup T_k)$ is finite since $S_k \cup T_k$ is $A_i \cup A_j$ -dispersed, and non-empty since (S_k, T_k) separates A_i from A_j , with in particular $E(W) \cap T_k \neq \emptyset$ if $V(W) \cap S_k = \emptyset$. Thus, if $V(W) \cap S = \emptyset$, there is $k \in K$ such that $V(W) \cap S_k = \emptyset$, hence $E(W) \cap T_k \neq \emptyset$; and by the finiteness of $E(W) \cap T_k$, this implies that $E(W) \cap T \neq \emptyset$. Therefore $(S, T) \in \text{SEP}(\mathcal{A})$. \square

3. \mathcal{A} -paths

Definition 3.1. Let τ_0 and τ_1 be two distinct ends of G . A $\tau_0\tau_1$ -path is a finite or infinite path (ray or double ray) W of G such that:

- if W is a double ray then $W = R_0 \cup R_1$ with $R_i \in \tau_i$ for $i = 0, 1$;
- if W is a ray $\langle x_0, x_1, \dots \rangle$ then $W \in \tau_i$ for some $i \in \{0, 1\}$ and $x_0 \in V_{\tau_{1-i}}$;
- if W is a path $\langle x_0, \dots, x_n \rangle$ then $x_0 \in V_{\tau_i}$ and $x_n \in V_{\tau_{1-i}}$ for some $i \in \{0, 1\}$.

3.2. For a set $\mathcal{A} := \{A_i : i \in I\}$ of pairwise disjoint subsets of $\mathfrak{X}(G)$, we call a $\tau_i\tau_j$ -path an \mathcal{A} -path if $(\tau_i, \tau_j) \in A_i \times A_j$ and $\{i, j\} \in [I]^2$, and we denote by $\mathcal{W}_{\mathcal{A}}$ the set of \mathcal{A} -paths, and by $\text{DISJ}(\mathcal{A})$ the set of all sets of pairwise disjoint \mathcal{A} -paths. Finally we set

$$\text{disj}(\mathcal{A}) := \sup\{|\mathcal{D}| : \mathcal{D} \in \text{DISJ}(\mathcal{A})\}.$$

Theorem 3.3. *If \mathcal{A} is such that $A_i \cap \overline{\bigcup_{j \neq i} A_j} = \emptyset$ for every $i \in I$, then there is a set of pairwise disjoint \mathcal{A} -paths of cardinality $\text{disj}(\mathcal{A})$. Furthermore, if $\text{disj}(\mathcal{A})$ is infinite, then every maximal element of $\text{DISJ}(\mathcal{A})$ with respect to inclusion, is of maximum cardinality.*

Proof. This is obvious when $\text{disj}(\mathcal{A})$ is finite, and when it is uncountable since every \mathcal{A} -path is countable.

Suppose that $\text{disj}(\mathcal{A}) = \omega$, and let \mathcal{D} be a maximal element of $\text{DISJ}(\mathcal{A})$. Assume $|\mathcal{D}| < \omega$. Then, since $\text{disj}(\mathcal{A}) = \omega$ and since \mathcal{D} is maximal with respect to inclusion, \mathcal{D} has an infinite element containing a ray R such that, for any positive integer n , there is a $\mathcal{D}_n \in \text{DISJ}(\mathcal{A})$ with $|\mathcal{D}_n| \geq n$ and $W \cap R \neq \emptyset$ for all $W \in \mathcal{D}_n$. Thus $[R]_G \in A_i \cap \overline{\bigcup_{j \neq i} A_j}$ for some $i \in I$, a contradiction with the hypothesis. Hence $|\mathcal{D}| = \omega$. \square

Using the extension Theorem 1.2 of Mader's Theorem, we will now prove the main result, which will generalize Theorem 3.4 of [8].

Theorem 3.4. *If $\mathcal{A} := \{A_i : i \in I\}$ is such that $A_i \cap \overline{\bigcup_{j \neq i} A_j} = \emptyset$ for every $i \in I$, then $\text{disj}(\mathcal{A}) = \text{sep}(\mathcal{A})$.*

Proof. (a) Let (S, T) be an \mathcal{A} -separator, and W be an \mathcal{A} -path. Suppose that $V(W) \cap S = \emptyset$ and $|V(W) \cap \mathcal{B}(X, G - S)| < 2$ for any component X of $G(T)$. Then, by condition (iii) of Definition 2.1, W is not a double ray. Thus it has at least an endpoint x which belongs to V_τ for some $\tau \in \bigcup \mathcal{A}$ such that, by the definitions of V_τ and of an \mathcal{A} -separator, there is an infinite set of pairwise internally disjoint rays in τ originating at x and meeting S or T . But this implies that $S \cup \bigcup T$ has an infinite subset which is concentrated in τ , a contradiction with condition (ii) of Definition 2.1. Hence $\text{disj}(\mathcal{A}) \leq \text{sep}(\mathcal{A})$.

(b) By Theorem 2.3, (2.2.2) and 0.6, there exists a finitary \mathcal{A} -separator S . Let $\Gamma := \{X \in \mathcal{C}_{G-S} : \mathfrak{T}_X(G) \cap \bigcup \mathcal{A} \neq \emptyset\}$, and let $X \in \Gamma$. Since S is finitary, the set $S_X := \mathcal{B}(G \mid S, X)$ is finite; and since S is an \mathcal{A} -separator, $A_X := \bigcup \mathcal{A} \cap \mathfrak{T}_X(G) \subseteq A_i$ for some $i \in I$. Let F_X be a finite subset of $S_X \cup V(X)$ such that $S_X \cap V(\mathcal{C}_{G-F_X}(A_X)) = \emptyset$, and which is of minimum cardinality. By Theorem 4.7 of [5] $|F_X|$ is equal to the maximum number of pairwise disjoint (S_X, A_X) -paths (a (S_X, A_X) -path is a path or a ray W originating in S_X such that, if W is a ray then $[W]_G \in A_X$, and otherwise $V(W) \cap V_{A_X} \neq \emptyset$, and which is minimal with this property with respect to inclusion).

For $i \in I$, we denote by Γ_i the subset of Γ such that $A_i = \bigcup_{X \in \Gamma_i} A_X$, and by a_i a vertex so that $a_i \notin V(G)$, and $a_i \neq a_j$ if $i \neq j$. Finally we define the graphs:

$$G'_{\mathcal{A}} := G - \bigcup_{X \in \Gamma} \mathcal{C}_{G-F_X}(A_X)$$

and

$$G_{\mathcal{A}} := \bigcup_{1 \leq i \leq p} \left\{ \langle a_i, x \rangle : x \in \bigcup_{X \in \Gamma_i} F_X \right\} \cup G'_A.$$

Notice that, by the definitions of S and of the F_X 's $V_{\tau_i} \cap V_{\tau_j} \subseteq V(G_{\mathcal{A}})$ for all $\{i, j\} \in [I]^2$ and $(\tau_i, \tau_j) \in A_i \times A_j$.

Then, still by the definition of the F_X 's, for any family of pairwise disjoint \mathcal{A} -paths \mathcal{D} there is a family of pairwise internally disjoint $\{a_i : i \in I\}$ -paths \mathcal{D}' such that

$$\{W \cap G'_{\mathcal{A}} : W \in \mathcal{D}\} = \{W \cap G'_{\mathcal{A}} : W \in \mathcal{D}'\};$$

and conversely. Therefore $\text{disj}(\mathcal{A}) = \text{disj}(\{a_i : i \in I\})$. On the other hand

$$\text{SEP}_{G_{\mathcal{A}}}(\{a_i : i \in I\}) = \{(S, T) \in \text{SEP}(\mathcal{A}) : S \cup T \subseteq V(G'_{\mathcal{A}})\}.$$

Consequently, by the extension Theorem 1.2 of Mader's Theorem,

$$\text{sep}(\mathcal{A}) \geq \text{disj}(\mathcal{A}) = \text{disj}_{G_{\mathcal{A}}}(\{a_i : i \in I\}) = \text{sep}_{G_{\mathcal{A}}}(\{a_i : i \in I\}) \geq \text{sep}(\mathcal{A}).$$

Hence $\text{disj}(\mathcal{A}) = \text{sep}(\mathcal{A})$. \square

When \mathcal{A} has only two elements, Theorem 3.3 of [8] gives the following precision: *If $|\mathcal{A}| = 2$ then there is $(S, \emptyset) \in \text{SEP}(\mathcal{A})$ such that $|S| = \text{sep}(\mathcal{A}) = \text{disj}(\mathcal{A})$.*

References

- [1] H. Freudenthal, Über die Enden diskreter Räume und Gruppen, *Comment. Math. Helv.* 17 (1944) 1–38.
- [2] R. Halin, Über unendliche Wege in Graphen, *Math. Ann.* 157 (1964) 125–137.
- [3] R. Hallin, A note on Menger's Theorem for infinite locally finite graphs, *Abh. Math. Sem. Univ. Hamburg* 40 (1974) 111–114.
- [4] W. Mader, Über die Maximalzahl kreuzungsfreier H -Wege. *Arch. Math.* 31 (1978) 387–402.
- [5] N. Polat, Aspects topologiques de la séparation dans les graphes infinis I, *Math. Z.* 165 (1979) 73–100.
- [6] N. Polat, Aspects topologiques de la séparation dans les graphes infinis II, *Math. Z.* 165 (1979) 171–191.
- [7] N. Polat, Développements terminaux des graphes infinis I, Arbres maximaux coterminaux. *Math. Nachr.* 107 (1982) 283–314.
- [8] N. Polat, A Mengerian theorem for infinite graphs with ideal points, *J. Combin. Theory Ser. B* 51 (1991) 248–255.
- [9] R. Rado, A selection lemma, *J. Combin. Theory* 10 (1971) 176–177.
- [10] B. Zelinka, Uneigentliche Knotenpunkte eines Graphen, *Časopis Pěst. Math.* 95 (1970) 155–169.